

Precision charmonium and D physics from lattice QCD and determination of the charm quark mass

C. T. H. Davies (for the HPQCD collaboration)

Department of Physics and Astronomy, University of Glasgow, Glasgow G12 8QQ, UK

I will describe recent results from the HPQCD collaboration using a new very accurate method for charm quarks in lattice QCD, that we have used in calculations including the full effect of u , d and s sea quarks. Multiple values of the lattice spacing and of the u , d and s sea quark masses allow us to extrapolate reliably, with a full error budget, to the real world. This opens up the field of charm physics to precision lattice QCD tests. So far we have calculated the D and D_s meson masses to 6 MeV, having fixed the charm quark mass from the η_c meson. Our D and D_s decay constants (determined to 2%) make an interesting comparison to CLEO-c results as we await improved experimental errors. We are also able to determine the charm quark mass to an accuracy of 1% using charmonium correlators and high-order continuum QCD perturbation theory. Future calculations are briefly discussed.

1. INTRODUCTION

Lattice QCD is now established as a precision tool for ‘gold-plated’ hadron physics [1], enabling us both to test QCD to high accuracy but also to use it to determine parameters of the Standard Model such as quark masses and CKM matrix elements.

Figure 1 shows the current status of HPQCD lattice calculations of the masses of gold-plated mesons. These are mesons which are far from strong decay thresholds and whose masses can then be accurately determined both theoretically and experimentally. The lattice calculations are done including the full effect of sea u , d and s quarks (using the MILC collaboration gluon configurations) and at multiple values of the lattice spacing so that systematic errors from working on a space-time lattice can be removed as far as possible, and a full error budget worked out. A lot of the gold-plated mesons contain valence c or b quarks. Whilst work continues on Υ , B and related mesons there has been a lot of recent progress on mesons containing c quarks, and I report on that here. One of the key issues in charm physics is to test lattice QCD against experiment as a precursor to using accurate lattice QCD results for B physics along with experiment to determine key CKM elements. We now have a 2% accurate lattice QCD calculation of the decay constants of the D and D_s mesons [2] and the recent experimental updates on this from CLEO-c have thrown up an interesting picture. This is discussed in section 2. Further theoretical tests of our charm methods have led to a 1% accurate determination of the charm quark mass, using methods pioneered in continuum QCD [3], and this is discussed in section 3.

2. CHARM QUARKS IN LATTICE QCD

In continuum QCD both charm and bottom quarks are described as heavy because their masses are significantly greater than the QCD scale, Λ_{QCD} , of a few hundred MeV. Special techniques can be applied to heavy quarks because they are nonrelativistic in their bound states and $\alpha_s(m_Q)$ is relatively small. This is important information that can be applied to handling b quarks in lattice QCD also. For c quarks it turns out not to be nearly so useful because the charm quark mass in lattice units, $m_c a$, is not that large for values of the lattice spacing currently in use, and will get even smaller as we make finer and finer lattices. We believe that a better strategy for c quarks is to treat them in the same way as light u , d and s quarks. This allows us to make use of light quark symmetries of the (lattice) QCD Lagrangian so that, for example, the annihilation rate of pseudoscalar mesons can be directly calculated without the need for a renormalisation constant. We can also apply standard variance reduction techniques and find then that lattice statistical errors for D/D_s mesons are as small (less than 1%) as those for π and K .




Figure 1: The gold-plated meson spectrum from lattice QCD. Lines represent experimental values. Open circles are lattice results, distinguishing those hadron masses used to fix the quark mass parameters and those which were predictions ahead of experiment [4, 5].

The worry, however, with c quarks, is that $m_c a$ is still quite large and this means that $m_c a$ will set the scale for discretisation errors that appear as a result of having a finite lattice spacing. Although these can be extrapolated away with multiple values of the lattice spacing, a large extrapolation will lead to a large resulting error. Typically we would expect, for some quantity m , to obtain a result at non-zero lattice spacing, a , that behaves as:

$$m = m_{a=0}(1 + A(m_c a)^2 + B(m_c a)^4 + \dots). \quad (1)$$

For $a \approx 0.1\text{fm}$, $(m_c a) \approx 0.4$ and $(m_c a)^2$ terms could lead to a 20% error. These are present in most light quark actions (where they do not cause a big problem). We designed an action in which these terms were not present so that errors of only a few percent would be made even for c quarks, and once extrapolations were made to zero lattice spacings any remaining errors would be very small. The action is based on the improved staggered quark action used for sea quarks in the MILC configurations, and is called the Highly Improved Staggered Quark action (HISQ) [6].

2.1. Results for D and D_s mesons

A lattice calculation proceeds by generating sets of gluon fields that are ‘typical snapshots of the vacuum’. The MILC collaboration now have many such sets of gluon fields at 5 different values of the lattice spacing including the full effect of u , d and s sea quarks using the improved staggered quark formalism [7]. We have used c and light valence HISQ quarks on these gluon configurations to calculate correlators for η_c , D , D_s , K and π mesons. By studying the correlators as a function of lattice time in the large time limit, we can then calculate the masses of these mesons and the decay amplitude known as the decay constant. This is defined in pure QCD as f_H where $f_H m_H = \langle 0 | \bar{\psi} \gamma_0 \gamma_5 \psi | H \rangle$, and is the probability that the quark and antiquark are in the same place to annihilate




Figure 2: A comparison of lattice results for the D (leftmost points) and D_s (rightmost points) decay constants with experiment. The experimental results are obtained from the leptonic decay rate using CKM elements V_{cs} and V_{cd} from elsewhere [8, 9, 10]. CLEO-c presented an update at this conference [11]. HPQCD results are from [2] using HISQ quarks. The Fermilab/MILC results are an update this year [12], ETMC results are new [13] and do not include the s sea quarks that are present in the real world. Neither of these results use a formalism as accurate as HISQ and hence have larger errors. Currently there is agreement between lattice and experiment for f_D but not for f_{D_s} , when comparing HPQCD and CLEO-c results.

to a W boson. When calculated in QCD, it takes into account all the strong interactions that keep the quark and antiquark bound into the meson.

The quark masses need to be fixed in any QCD calculation and we do this by adjusting them until the η_c , K and π meson masses are correct. The last two require extrapolation in the u/d quark mass (the sea and valence masses are chosen to be the same) using chiral perturbation theory because the lattice calculation has to be done at u/d masses that are larger than the physical values. D and D_s masses have no further free parameters and we obtain values for them in agreement with experiment and with 6 MeV errors (again after extrapolation in the u/d mass to the physical point) [2]. This is a very strong test both of our control of discretisation errors and of QCD itself because it shows that charmonium and charm-light systems, with very different dynamics, are simultaneously described a single QCD Lagrangian.

The rate for annihilation to a $W(\rightarrow l\bar{\nu})$ is proportional to $f_H^2 V_{ab}^2$ where V_{ab} is the appropriate CKM element. Our results for f_K/f_π mesons can be used, with experimental rates, to determine V_{us} to 0.6% [2]. Our results for D and D_s decay constants have 2% errors and can be compared to experimental determinations obtained by dividing the leptonic rate by known values of V_{cs} and V_{cd} . Following our calculations the results from the experiment have become very exciting this year, as shown in Figure 2. A new result for f_D from CLEO-c [10] agrees well with our result, but values for f_{D_s} do not. The discrepancy there amounts to 3σ where σ comes from experiment because our error is so small. This has led to speculation of new physics [14], since the disagreement is a serious one and is the only such disagreement from the 15 or so quantities that have now been accurately calculated in lattice QCD and compared to experiment, see Figure 1. Final results from CLEO-c on f_{D_s} should reduce the error bars further and either confirm, or not, this hypothesis.

Meanwhile, we have continued to test our control of lattice systematic errors by calculating vector (e.g. J/ψ) electromagnetic decay rates and hyperfine splittings such as $J/\psi - \eta_c$ and $D_s^* - D_s$. These are in good agreement with experiment and we are finalising error budgets for them. Future work aims at few percent errors for semileptonic

form factors for $D \rightarrow K, \pi$ for comparison with experiment.

3. THE CHARM QUARK MASS

We have developed, in collaboration with continuum QCD theorists, a new method for determining heavy quark masses. So far we have applied this to determine m_c to 1% and the agreement between our result for the mass in the $\overline{MS}$ scheme at 3 GeV, 0.986(10) GeV [15], and the result obtained by analogous continuum techniques, 0.986(13) GeV [16], is another strong test of our control of systematic errors in handling charm quarks in lattice QCD.

Lattice QCD calculations encompass energy scales from low-energies and long-distances (which we use to extract hadron masses) to relatively high-energies set by the inverse of the lattice spacing. These high-energies are well into the perturbative regime and we make use of this, for example, in the accurate determination of α_s [17]. Our charmonium (η_c or J/ψ) correlators are also perturbative at short lattice times, before the ground state meson dominates (and we extract the ground state meson mass). We can access this perturbative region by taking 'time-moments', $G_n = \sum_t (t/a)^n G(t)$, of the correlator, $G(t)$. Extrapolation of these to the continuum allows them to be compared to continuum QCD perturbation theory which should be accurate for small enough values of n and is known in some cases up to and including α_s^3 terms. The analogous continuum calculation [16] effectively determines correlator moments in the vector case from $R(e^+e^- \rightarrow \text{hadrons})$ in the charm region. Our calculation in the vector (J/ψ) case agrees with experiment but our most accurate result, quoted above, comes from the pseudoscalar (η_c) [15].

The determination of m_c from the more conventional lattice method of converting the bare lattice mass to the $\overline{MS}$ scheme using lattice QCD perturbation theory, agrees well with this result [18]. The accurate results for m_c can be leveraged into an accurate result for light quark masses by determining ratios such as m_c/m_s from which renormalisation constants between the lattice and the continuum cancel [19].

Acknowledgments

I am grateful to my colleagues in the HPQCD collaboration, and to members of the Fermilab lattice and MILC collaborations, for many useful discussions. Work supported by the STFC and the Leverhulme Trust.

References

- [1] C.T. H. Davies *et al*, Fermilab lattice/HPQCD/MILC, Phys. Rev. Lett.**92**:022001 (2004) [hep-lat/0304004].
- [2] E. Follana *et al*, HPQCD collaboration, Phys. Rev. Lett.**100**:062002 (2008) arXiv:0706.1726[hep-lat].
- [3] J. H. Kühn, this Proceedings, arXiv:0809.1780[hep-ph].
- [4] A. Gray *et al*, HPQCD collaboration, Phys. Rev. D**72**:094507 (2005) [hep-lat/0507013].
- [5] I. F. Allison *et al*, HPQCD/Fermilab Lattice collaborations, Phys. Rev. Lett**94**:172001 (2005) [hep-lat/0411027].
- [6] E. Follana *et al*, HPQCD collaboration, Phys. Rev. D**75**:054502 (2007) [hep-lat/0610092].
- [7] C. Bernard *et al*, MILC collaboration, PoS LAT2007:090 2007 arXiv:0710.1118[hep-lat].
- [8] L. Widhalm, Belle collaboration, Proceedings of Charm 2007, arXiv:0710.0420.
- [9] B. Aubert *et al*, BaBar collaboration, Phys. Rev. Lett.**98**:141801 (2007) [hep-ex/0607094].
- [10] K. Ecklund *et al*, CLEO-c, Phys. Rev. Lett**100**:161801 (2008) arXiv:0712.1175[hep-ex]; B. Eisenstein *et al*, CLEO-c, arXiv:0806.2112[hep-ex];
- [11] L. Zhang, CLEO-c, this Proceedings, arXiv:0810.2328[hep-ex].
- [12] P. Mackenzie *et al*, Fermilab Lattice, MILC collaborations, Proceedings of LAT08.
- [13] B. Blossier *et al*, ETMC collaboration, Proceedings of LAT08, arXiv:0810.3145.
- [14] See, for example, Dobrescu and Kronfeld, Phys. Rev. Lett.**100**:241802 (2008) arXiv:0803.0512[hep-ph].

- [15] I. Allison *et al*, HPQCD collaboration + K. G. Chetyrkin, J. H. Kühn, M. Steinhauser and C. Sturm, arXiv:0805.2999[hep-lat], Phys Rev. D (in press).
- [16] J. H. Kühn, M. Steinhauser, C. Sturm, Nucl. Phys. B**778**:192 (2007) [hep-ph/0702103].
- [17] C. T. H. Davies *et al*, HPQCD collaboration, arXiv:0807.1687[hep-lat].
- [18] I. F. Allison *et al*, HPQCD collaboration, Proceedings of LAT08, arXiv:0810.0285[hep-lat].
- [19] C. T. H. Davies, G. P. Lepage *et al*, HPQCD collaboration, Proceedings of LAT08, arXiv:0810.3548[hep-lat].